

THE SPECTRUM OF THE DIFFUSE GALACTIC LIGHT I: THE MILKY WAY IN SCATTERED LIGHT

TIMOTHY D. BRANDT AND B. T. DRAINE¹*Draft version November 18, 2011*

ABSTRACT

We measure the optical spectrum of the Diffuse Galactic Light (DGL) – the local Milky Way in reflection – using 92,000 blank-sky spectra from the Sloan Digital Sky Survey (SDSS). We correlate the SDSS optical intensity in regions of blank sky against 100 μm intensity independently measured by the COsmic Background Explorer (COBE) and InfraRed Astronomy Satellite (IRAS) satellites, which provides a measure of the dust column density times the intensity of illuminating starlight. The spectrum of scattered light is very blue and shows a clear 4000 $\text{\AA}$ break and broad Mg b absorption. This is consistent with scattered starlight, and the continuum of the DGL is well-reproduced by a simple radiative transfer model of the Galaxy. We also detect line emission in H α , H β , [N II], and [S II], consistent with scattered light from the local interstellar medium (ISM). The strength of [N II] and [S II], combined with upper limits on [O III] and [He I], indicate a relatively soft ionizing spectrum. We find that our measurements of the DGL can constrain dust models, favoring a grain size distribution with relatively few large grains. We also estimate the fraction of high-latitude H α which is scattered to be $19 \pm 4\%$.

Subject headings: ISM: dust, extinction; Scattering; Methods: statistical

1. INTRODUCTION

All astronomical observations include light from diffuse sources other than the target object. In ground-based data, the dominant sources of contamination in the optical are airglow, scattered sunlight, and artificial sources. Space-based missions must still contend with zodiacal light, and all observations will include some emission and scattering from the Galaxy’s interstellar medium (ISM) – the diffuse Galactic light (DGL).

The first quantitative measurements of the DGL were photoelectric measurements by Elvey & Roach (1937) at $\lambda \approx 4500 \text{ \AA}$; after subtracting the zodiacal and airglow contributions, the DGL was detected for $|b| \lesssim 35^\circ$. Subsequent studies from the ground (Elsässer & Haug 1960) and a sounding rocket (Wolstencroft & Rose 1966) found intensities at $|b| \approx 5^\circ$ corresponding to ~ 50 10th magnitude stars per square degree, or $\lambda I_\lambda \approx 5 \times 10^{-4} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$. Subsequent observations from satellites (Lillie & Witt 1976; Morgan et al. 1978; Henry 1981; Zvereva et al. 1982; Martin et al. 1990; Murthy et al. 1990; Hurwitz et al. 1991; Murthy et al. 1991; Sasseen & Deharveng 1996; Seon et al. 2011) extended these studies into the vacuum ultraviolet (UV). Most of these studies were broadband, but Martin et al. (1990) detected fluorescent emission from UV-pumped H₂.

The spectrum of the DGL contains a wealth of information about the physical environment where it originates and the dust that emits or scatters it into our line-of-sight. In this paper, we present a novel way of measuring the spectrum of light scattered by the Galactic ISM. This is a spectrum of the Galaxy in reflection, plus possible luminescence from interstellar dust.

2. METHODOLOGY

The optical surface brightness of the DGL, $\lambda I_\lambda \approx 10^{-4} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$, is far too low to measure a spectrum directly. Further, any such spectrum would be a combination of terrestrial airglow, scattered artificial light, zodiacal light, scattering and emission by interstellar dust, emission by diffuse gas, and unresolved background objects. In this section we describe a novel technique to measure the spectrum of scattering by the Galactic ISM. We use 92,000 sky spectra from the Sloan Digital Sky Survey (SDSS, York et al. 2000), correlating their intensities against independently measured 100 μm emission to isolate the components of the DGL associated with interstellar dust. In a companion paper (Brandt & Draine 2011, in preparation) we use the full-sky H α map compiled by Finkbeiner (2003) to isolate the component of the DGL associated with emission by diffuse H II.

2.1. The SDSS Sky Fibers

The Seventh Data Release of the SDSS (Abazajian et al. 2009) contains more than 1.6 million spectra of stars, galaxies, and quasars, making it by far the largest such dataset ever assembled. The spectra were taken by plugging 640 fibers into a $\sim 7 \text{ deg}^2$ plate, with each 2.96'' fiber feeding the light of its target object into a pair of spectrographs. Each group of 640 spectra was then calibrated and sky-subtracted by the SDSS spectroscopic pipeline (Stoughton et al. 2002; Burles & Schlegel unpub.).

To obtain an accurate measurement of the sky background, a minimum of 32 fibers on each plate were placed on blank sky regions. These positions were relatively uniformly distributed over each plate, and required no detection by the photometric pipeline to 5σ in any band (Lupton, priv. commun.). A few ($\sim 1\%$) of the sky fibers were erroneously placed over bright sources, but these were flagged and removed by the reduction pipeline, leaving a total of about 92,000 blank sky spectra used to compute the background flux. The sky fibers were used to con-

¹ Department of Astrophysical Sciences, Peyton Hall, Princeton University, Princeton, NJ 08544, USA

struct a “supersky” spectrum for each plate scaled to unit airmass. This spectrum was then rescaled to the airmass at each fiber (including the sky fibers themselves) and subtracted from that fiber’s spectrum. Each plate also included 8 F dwarfs as spectrophotometric standards, 8 F subdwarfs as reddening standards, and 2 hot subdwarfs (Stoughton et al. 2002). The residual sky spectra, along with all of the other spectra, were flux-calibrated to these standards.

The sky spectra on a given plate show modest fiber-to-fiber variation, but are dominated by noise associated with terrestrial airglow. Hidden within this noise are real variations due to extraterrestrial sources: zodiacal light, scattering and emission by diffuse interstellar dust, emission by diffuse gas, and faint, unresolved background sources. We can isolate the components associated with Galactic dust using the independently measured InfraRed Astronomy Satellite (IRAS) 100 μm map, reduced and calibrated by Schlegel et al. (1998), hereafter SFD. The bulk of this emission comes from thermally radiating dust grains heated to ~ 18 K by starlight. By tracing interstellar dust illuminated by Milky Way stars, the 100 μm map allows us to measure the spectrum of the Galaxy in scattered light.

2.2. Correlating Against 100 μm Intensity

In 1983, IRAS mapped the entire sky at 12, 25, 60, and 100 microns (Neugebauer et al. 1984). SFD later smoothed the 100 μm map and corrected it for point sources and zodiacal light, creating a map of diffuse Galactic infrared emission. They further used COsmic Background Explorer (COBE) data (Boggess et al. 1992) at 240 μm to estimate the temperature of the dust, ultimately producing a $\sim 6'$ resolution map of 100 μm emission and a lower resolution temperature map suitable to convert 100 μm emission into a dust column density. This map was intended mainly to estimate and correct for Galactic extinction. Here, we use it to correlate illuminated dust with residual intensity in the SDSS sky fibers.

In the optically thin limit, we expect the intensity of scattered starlight to be proportional to the column density of dust times the intensity of the illuminating starlight. For dust with opacity $\propto \nu^\beta$ near 100 μm , the intensity of the illuminating (and hence, scattered) starlight should be proportional to $T_{\text{dust}}^{4+\beta}$. Planck Collaboration et al. (2011) have found that the $\lambda > 100$ μm emission is well-approximated by dust with $\beta \approx 1.8$. We therefore expect the SDSS sky fiber residual intensity to be roughly proportional to $\tau_{100\mu\text{m}} T_{\text{dust}}^{5.8}$, where the optical depth at 100 μm , $\tau_{100\mu\text{m}}$, is proportional to the dust column density. We expect the 100 μm intensity itself to have a temperature dependence. For 18 K dust radiating at 100 μm , $h\nu/k_B T \approx 8 \gg 1$, so that

$$\left(\frac{d \ln I_\nu}{d \ln T} \right) \bigg|_{\substack{\lambda=100\mu\text{m} \\ T=18\text{K}}} \approx \frac{h\nu}{k_B T} \approx 8. \quad (1)$$

We use the measured 100 μm intensity to trace the product of the intensity of the illuminating starlight and the dust column. In practice, temperature variations are sufficiently small that using the SFD column density times $T^{5.8}$ produces results indistinguishable from those simply

using 100 μm intensity.

Two complications prevent us from assuming a linear relationship between sky fiber residuals and 100 μm intensity:

1. Such a model neglects self-absorption by optically thick dust.
2. The spectroscopic pipeline has already subtracted a scaled sky spectrum from each fiber, which includes a component of the DGL. This component differs from plate to plate.

We avoid the first problem by excluding spectra and entire plates where the dust is optically thick to visible light, with $A_V \gtrsim 0.5$ according to SFD; our results are insensitive to the precise value of this threshold. For a dust temperature of 18 K, this corresponds to 100 μm emission exceeding ~ 10 MJy sr $^{-1}$. We address the second point by assuming residual optical intensity to be proportional to the excess 100 μm intensity *relative to the average over that fiber’s plate*. Our model is then

$$\lambda I_{\lambda, \text{sky}, j, p} = \alpha_\lambda [(\nu I_\nu)_{100\mu\text{m}, j, p} - \langle (\nu I_\nu)_{100\mu\text{m}} \rangle_p], \quad (2)$$

where $I_{\lambda, \text{sky}, j, p}$ is the residual intensity in sky fiber j on plate p at wavelength λ , $(I_\nu)_{100\mu\text{m}, j, p}$ is the 100 μm intensity at fiber j ’s location, $\langle \rangle_p$ denotes an average over the sky fibers on plate p , and α_λ is a dimensionless number that describes the relative strength of scattered and thermal emission. We then solve for the best-fit spectrum of coefficients α_λ , our *correlation spectrum*. Defining

$$y_{\lambda, j, p} \equiv \lambda I_{\lambda, \text{sky}, j, p} \quad \text{and} \quad (3)$$

$$x_{j, p} \equiv (\nu I_\nu)_{100\mu\text{m}, j, p} - \langle (\nu I_\nu)_{100\mu\text{m}} \rangle_p, \quad (4)$$

the maximum likelihood estimate for α_λ is

$$\alpha_\lambda = \left(\sum_{j, p} \frac{(y_{\lambda, j, p}) (x_{j, p})}{\sigma_{\lambda, y, j, p}^2} \right) \left(\sum_{j, p} \frac{x_{j, p}^2}{\sigma_{\lambda, y, j, p}^2} \right)^{-1}, \quad (5)$$

and its variance is

$$\sigma_\lambda^2 = \left(\sum_{j, p} \frac{x_{j, p}^2}{\sigma_{\lambda, y, j, p}^2} \right)^{-1}. \quad (6)$$

In Equations (5) and (6), $\sigma_{\lambda, y, j, p}^2$ is the variance of $y_{\lambda, j, p}$ as estimated by the SDSS pipeline.

By using the residual, sky-subtracted intensity, our model adopts the flux calibrations performed by the SDSS spectroscopic pipeline. Beginning with the Sixth Data Release, SDSS spectra have been flux-calibrated to PSF magnitudes appropriate for point sources rather than fiber magnitudes appropriate for extended sources. Dividing by the SDSS fiber aperture therefore gives an incorrect intensity. Figure 4 of Adelman-McCarthy et al. (2008) shows the difference between PSF and fiber magnitudes of point sources to be very nearly Gaussian with a mean of 0.35 magnitudes. The typical seeing for spectroscopy was poor ($\sim 2''$); as a result, the aperture correction was found to be a very weak function of wavelength. We therefore recalibrate all of our spectra to fiber magnitudes using the average flux conversion factor of $10^{0.35/2.5} = 1.38$ before dividing by the $2.96''$ fiber aperture.

We would like to apply the model in Equations (2)–(6) to each wavelength observed by SDSS. However, the wavelengths that were observed differed slightly from night to night. This variation is only a few tenths of a percent, but is sufficient to blur spectral features if not removed. We therefore define a new wavelength array of 4000 elements, similar to but slightly larger than the 3852 element array in most SDSS spectra. We use cubic splines to interpolate all spectra and errors onto this new array. The interpolation introduces a correlation between neighboring wavelength elements. In Section 4.1, we discuss this in detail, and demonstrate that our measured spectra and errors are statistically very well-behaved.

Figure 1 illustrates our model for the spectrum of scattered light. To increase the signal-to-noise ratio in Figure 1, we have binned each spectrum’s ~ 60 wavelength elements from 6900–7000 Å. Each 2.96'' patch of blank sky thus contributes a single point: its residual intensity averaged over the interval from 6900 to 7000 Å. After discarding bad sky fibers flagged by the spectroscopic pipeline along with fibers and plates with $(I_\nu)_{100\mu\text{m}} > 10 \text{ MJy sr}^{-1}$ (corresponding to $A_V \gtrsim 0.5$), we are left with nearly 90,000 points representing over 5 million intensity measurements. We indicate these points by logarithmically spaced contours where their density is extremely high. We then fit them with Equation (2), correlating the residual optical intensities against their fibers’ excess 100 μm emission. There is a positive correlation between the two quantities ($\alpha_\lambda > 0$), driven by the entire ensemble of points and significant at more than 70σ .

The best-fit slope in Figure 1 is biased low by its neglect of errors in the 100 μm map and by structure unresolved by IRAS, but by a factor independent of wavelength. Chromatic effects due to seeing and atmospheric dispersion are small (Adelman-McCarthy et al. 2008). We discuss the calibration of the correlation spectrum in Sections 4.2 and 4.3.

2.3. Sky Coverage

The SDSS focused much of its spectroscopy on the Northern Galactic Cap because of that region’s low levels of extinction. For the same reason, many of its plates are not very useful for inferring the spectrum of scattered light. The part of the SDSS footprint that is most useful for our purposes has significant small-scale variation in 100 μm emission, and is disproportionately in regions of low Galactic latitude.

Figure 2 shows the sky coverage of the SDSS spectroscopic survey (upper panel), and the sky coverage weighted by its power to affect the fit described by Equation (2) (lower panel). For any plate, the effect on the fit will be roughly proportional to the variance of the 100 μm intensity over that plate,

$$\sigma_{100\mu\text{m},p}^2 \equiv \langle I_{100\mu\text{m}}^2 \rangle_p - \langle I_{100\mu\text{m}} \rangle_p^2. \quad (7)$$

To obtain the weighted map, we have multiplied the fiber density by the ratio of $\sigma_{100\mu\text{m},p}^2$ to its average over the entire sample, $\langle \sigma_{100\mu\text{m}}^2 \rangle$.

Stripe 82 is clearly visible in the lower-left (and extreme lower-right) parts of each map. This region, defined by $-50^\circ < \text{RA} < 59^\circ$, $-1.25^\circ < \delta < 1.25^\circ$, was repeatedly imaged as part of the SDSS Supernova




FIG. 1.— Scatter plot showing the correlation between 100 μm intensity relative to the mean over an SDSS sky fiber’s plate and the residual sky intensity averaged over the interval from 6900 to 7000 Å (Equation 2). We use logarithmically spaced contours where the density of points is high. The data demand a non-zero slope, a component of the sky background associated with interstellar dust, at more than 70σ .

Survey (Adelman-McCarthy et al. 2006; Frieman et al. 2008; Abazajian et al. 2009). It was also the target of several “special” plates not part of the main SDSS surveys (Adelman-McCarthy et al. 2006). Stripe 82, along with a variety of patches at modest Galactic latitude and a swath at the southern limit of SDSS’s main footprint (in Galactic coordinates), contributes the bulk of our measured signal.

3. RESULTS: SPECTRA

Here we present our correlation spectra of the DGL, α_λ , computed as described in Section 2. We present three sets of spectra: the continuum of the scattered spectrum, and the spectra from 4830–5040 Å and from 6530–6770 Å to show emission lines. We also divide the sky into three regions by Galactic latitude and longitude to explore the spatial variation of the scattered light.

3.1. Continuum of the Correlation Spectrum

We obtain the spectrum of the DGL associated with 100 μm emission using the method described in Section 2.2. By fitting Equation (2) to the SDSS residual sky spectra, we obtain a dimensionless coefficient α_λ at each wavelength relating optical intensity to 100 μm emission. We then mask the nebular emission lines $\text{H}\alpha$, $\text{H}\beta$, $[\text{N II}] \lambda 6550$, $[\text{N II}] \lambda 6585$, $[\text{S II}] \lambda 6718$, and $[\text{S II}] \lambda 6733$ and bin the correlation spectra in intervals of 50 Å to increase the signal-to-noise ratio. We do not mask auroral lines like $[\text{O I}] \lambda 6300$, which are uncorrelated with the 100 μm intensity. As we show in Section 4.1, our errors are independent except for an interpolation effect, and they are normally distributed. The errors on the binned spectra pass the same statistical test.




FIG. 2.— Top panel: the sky coverage of the SDSS blank sky fibers in Galactic coordinates, centered around $(l, b) = (0, 0)$. Bottom panel: the sky coverage of the blank sky fibers weighted by $\sigma_{100\mu\text{m}, p}^2 / \langle \sigma_{100\mu\text{m}}^2 \rangle$ (Equation (7)), which provides an approximate measure of each fiber’s influence on the correlation spectrum. The bulk of the SDSS footprint is near the North Galactic Pole, with additional concentrated sampling in Stripe 82 near the South Galactic Pole. The useful survey footprint is much more heavily weighted toward equatorial regions.

Figure 3 shows the continuum spectrum of the DGL computed using all the sky fibers. The spectrum is very blue, yet it shows a clear 4000 Å break characteristic of old stellar populations. Broad Mg and Fe b absorption are also visible just blueward of 5200 Å. All of these characteristics are consistent with a continuum of scattered starlight. A simplified radiative transfer calculation, discussed in Section 5.1, confirms scattering as the source of the DGL and shows that it may be used to discriminate between dust models. The four plotted curves use two estimates of the continuum of the interstellar radiation field (ISRF) and two dust models, the Zubko et al. (2004) and Weingartner & Draine (2001) models (hereafter ZDA04 and WD01); an excess of large grains in the latter produces a redder scattered spectrum. As we discuss in Section 4.2, errors and small-scale structure in the 100 μm map bias our recovered correlation spectrum low by an unknown factor, which we estimate to be 2.1 ± 0.4 (see Section 4.2). We have therefore scaled the radiative transfer models to a common level to show their different shapes. The scale factors are indicated on the plot, and are all consistent with our estimate for the bias.

Figure 4 shows the continuum spectra from restricted areas of the sky. The left panel divides the sky by Galactic latitude, with the ranges chosen to have comparable 100 μm emission summed over our sky fibers. In the high-




FIG. 3.— The full-sky continuum of the scattered DGL, calculated as described in Section 2. We have masked nebular emission lines and binned the remaining spectrum into 50 Å intervals. The continuum is very blue and shows a clear 4000 Å break and Mg and Fe b absorption just blueward of 5200 Å; all of these features suggest scattered starlight. Simplified radiative transfer calculations, described in Section 5.1, confirm this and demonstrate that the DGL can discriminate between dust models. Because our correlation spectrum is biased low by an unknown factor (Section 4.2), we have scaled all spectra to a common level. The dashed lines use a featureless ISRF, while the solid lines use stellar population synthesis models. The ZDA04 model has fewer large grains and gives a better fit than WD01. We find no evidence of Extended Red Emission (ERE) around 7000 Å (Section 5.4).

est latitude bin, this flux is distributed over a much larger number of sky fibers and the resulting measurement is considerably noisier (see Figure 2). Still, the errors are reliable (see Section 4.1), and the shape differs from its values at lower latitude by many sigma. It is not clear whether the spatial variation is due to variation in the dust properties, in the illuminating starlight, systematic effects that become important when the signal is weak, or some combination of these factors. In particular, Yahata et al. (2007) find a correlation of SDSS galaxy density with SFD 100 μm emission in regions of low extinction, consistent with extragalactic contamination of the SFD map. Yahata et al. (2007) report that this contamination corresponds to an inferred $A_V \approx 0.01$, or $I_{100\mu\text{m}} \approx 0.2 \text{ MJy sr}^{-1}$. This would represent about 15% of the 100 μm intensity in our sky fibers with $|b| > 50^\circ$, and an even larger fraction of the intraplate variation to which we are sensitive.

The right panel of Figure 4 shows the continuum spectra for different regions in Galactic longitude. We take the longitude l to run from -180° to 180° , and choose our regions to be equal in area and symmetric about the Galactic center. About half of our signal comes from the region opposite the center, with $120^\circ < |l| < 180^\circ$. As with Galactic latitude, there are significant spatial variations in the correlation spectrum. It is unclear whether the redder spectrum in the direction of the Galactic cen-




FIG. 4.— The continuum spectra of the scattering component of the DGL for different regions of sky. These are identical to Figure 3 for restricted ranges of l and b ; note that we take $-180^\circ < l < 180^\circ$. The continuum shows some variation with Galactic latitude (left panel) and looks qualitatively different at the highest latitudes, potentially due to extragalactic contamination of the $100\ \mu\text{m}$ map (Yahata et al. 2007). The spectrum is also significantly redder toward the Galactic center (right panel). This spatial variation could be due to differences in the dust composition, in the illuminating starlight, or uncorrected systematic effects.

ter is due to larger grains, redder illuminating starlight, or some other effect. This spectrum is the most sensitive to the cutoff at high optical depth (Section 2.2), which may indicate significantly reddened illuminating starlight.

3.2. Emission Lines

Figure 5 shows the correlation spectrum in the wavelength ranges 4830–5040 Å and 6530–6770 Å, computed using all of the sky fibers without binning the α_λ . The spectrum of the DGL exhibits strong nebular emission lines which we masked to show the continuum in Figures 3 and 4. We detect H α , H β , [N II] λ 6550, [N II] λ 6585, [S II] λ 6718, and [S II] λ 6733 at high significance, but see only weak emission in the [O III] λ 5008 line excited by early O-type stars. Unfortunately, the [O II] λ 3727 line lies just outside the SDSS wavelength range. These emission lines likely represent scattered photons that were originally emitted from H II regions and the diffuse warm ionized medium (WIM).

We list line strengths in Table 1 measured as equivalent widths, which are unaffected by bias factors (Section 4.2). We use an interval of 6 SDSS wavelength elements, corresponding to a velocity range of $400\ \text{km s}^{-1}$, to measure line intensities. For the lines between 6500 and 6800 Å, we use the average intensity over the interval from 6600 to 6700 Å as our estimate of the continuum. For H β , we use 30 wavelength elements on each side of the line, running from 4829–4858 Å and from 4864–4893 Å. The ratio of the [N II] doublet, [N II] λ 6585 to [N II] λ 6550, provides a check on our recovered spectrum; our measured value of 2.8 ± 0.6 matches the ratio of 3 expected from the Einstein A coefficients.

In order to measure the strengths of the Balmer lines

in emission, we need to account for the fact that they appear in absorption in stellar spectra. Fortunately, the strength of the Balmer absorption lines in composite stellar spectra is closely correlated with that of the 4000 Å Calcium break. We measure the 4000 Å break in our continuum spectrum and use a range of model stellar spectra from Bruzual & Charlot (2003), hereafter BC03, to fit a linear relationship between the strength of the break and the Balmer equivalent widths. We use models of 6 Gyr of constant star formation, single stellar populations of 2.5 Gyr, 5 Gyr, and 11 Gyr, and two exponential star formation histories, each with metallicities of 0.02 and 0.008 ($Z_\odot$ and $0.4Z_\odot$); a combination of these models should provide a reasonable fit to stellar populations in the Solar neighborhood. Defining δ_{4000} to be the ratio of the integrated intensity between 3850 and 4000 Å to the integrated intensity between 4000 and 4150 Å, and computing the continua and the line widths as described above, we find best-fit relationships of

$$\left(\frac{\text{EW}(\text{H}_\beta)}{\text{\AA}}\right) \approx -2.2\delta_{4000} + 0.17 \quad \text{and} \quad (8)$$

$$\left(\frac{\text{EW}(\text{H}_\alpha)}{\text{\AA}}\right) \approx -1.5\delta_{4000} - 0.19 \quad (9)$$

for the model stellar spectra. The root-mean-square scatters of the BC03 equivalent widths around the fits given by Equations (8) and (9) are $\sim 0.06\ \text{\AA}$ and $\sim 0.1\ \text{\AA}$, respectively, significantly smaller than the errors in our measurements of the DGL. Correcting for stellar absorption increases our measured H α and H β line strengths by $\sim 20\%$.

The corrected equivalent width of H α in emission is




FIG. 5.— Emission lines in the DGL. $H\beta$, $H\alpha$, $[N II] \lambda 6550$, $[N II] \lambda 6585$, $[S II] \lambda 6718$, and $[S II] \lambda 6733$ are all detected with high significance, while we find little evidence of emission in $[O III] \lambda 5008$. The lack of $[O III]$ indicates a relatively soft spectrum of ionizing photons compared to average H II regions, while strong $[S II]$ and $[N II]$ indicate warm, $\sim 8,000$ K gas.

TABLE 1
EQUIVALENT WIDTHS AND LINE RATIOS IN THE DGL

Line	Equivalent Width [Å]	Energy Ratio of Line to	
		$H\alpha$	$H\beta$
$H\beta \lambda 4863$	4.8 ± 0.7^a	0.38 ± 0.05	1
$[O III] \lambda 4960$	1.6 ± 0.6	0.14 ± 0.05	0.38 ± 0.15
$[O III] \lambda 5008$	0.8 ± 0.6	0.07 ± 0.05	0.19 ± 0.15
$[He I] \lambda 5877$	0.3 ± 0.8	0.03 ± 0.07	0.09 ± 0.21
$[N II] \lambda 6550$	2.4 ± 0.5	0.19 ± 0.04	0.50 ± 0.12
$H\alpha \lambda 6565$	12.5 ± 0.5^a	1	2.64 ± 0.38
$[N II] \lambda 6585$	6.6 ± 0.5	0.53 ± 0.04	1.40 ± 0.22
$[S II] \lambda 6718$	5.7 ± 0.4	0.45 ± 0.04	1.20 ± 0.19
$[S II] \lambda 6733$	4.3 ± 0.4	0.34 ± 0.04	0.91 ± 0.15

^a Corrected for stellar absorption using Equations (8) and (9).

TABLE 2
THE EIGHT NEAREST O STARS^a

GOS ID	Other ID	Spectral Type	Distance (pc)
G006.28+23.5901	ζ Oph	O9.5V	112^{+3}_{-3}
G203.86−17.7401	δ Ori A	O9.5V	221^{+33}_{-25}
G206.45−16.5901	ζ Ori A	O9.7Ib	239^{+43}_{-32}
G202.94+02.2001	15 Mon	O7V	309^{+60}_{-43}
G255.98−04.7101	ζ Pup	O4I	335^{+12}_{-11}
G262.80−07.6901	γ^2 Vel	O9:I	349^{+44}_{-35}
G195.05−12.0001	λ Ori A	O8III	361^{+89}_{-60}
G206.82−17.3401	σ Ori AB	O9.5V	380^{+136}_{-87}

^a References: Maíz-Apellániz et al. (2004); Maíz Apellániz et al. (2008)

consistent with its value of 11 Å in the local ISRF (Draine 2011, Table 12.1), obtained by integrating the full-sky $H\alpha$ map compiled by Finkbeiner (2003) from the Wisconsin $H\alpha$ Mapper (WHAM, Reynolds et al. 2002), Southern $H\alpha$ Sky Survey Atlas (SHASSA, Gaustad et al. 2001), and Virginia Tech Spectral-Line Survey (VTSS, Dennison et al. 1998). The line ratios provide a probe of the average physical conditions in the local ISM. The strength of the singly ionized $[N II]$ and $[S II]$ lines and weakness of $[O III] \lambda 5008$ and $[He I] \lambda 5877$ indicate that most of the S, N, and O are singly ionized, while the He is largely neutral. This implies a lack of photons with $h\nu > 24.6$ eV in the Solar neighborhood, which can be understood from the lack of stars of spectral type O8 and earlier within 300 pc of the Sun. The nearest eight O stars are listed in Table 2. We discuss the physical conditions of the local ISM in more detail in Section 5.

Figure 6 shows the strength of the emission lines $H\alpha$, $[N II] \lambda 6550$, $[N II] \lambda 6585$, $[S II] \lambda 6718$, and $[S II] \lambda 6733$ relative to 100 μ m emission for different ranges of Galac-

tic longitude and latitude. The lines are somewhat stronger at low Galactic latitude and in the direction of the Galactic center, though this could reflect the relative number of nearby H II regions rather than the physical conditions in the ISM. An important caveat is that, because of spatial variations in the 100 μ m intensity, the correlation spectra in different regions of the sky need not share the same bias factor (Sections 4.2 and 4.3). The ratios of the $[S II]$ and $[N II]$ lines to $H\alpha$, which are robust to calibration difficulties, vary little across the sky.

4. CALIBRATION AND MEASUREMENT ERRORS

The standard errors on our correlation spectra α_λ are very nearly normally distributed with the variance given by our maximum likelihood estimator, up to a constant factor due to our interpolation of the original SDSS spectra onto a common wavelength array. The absolute calibration of the α_λ is more problematic. Our neglect of (unknown) measurement errors in the 100 μ m inten-




FIG. 6.— Emission line spectra of the dust component of the DGL as functions of Galactic longitude and latitude. The line ratios and equivalent widths depend only weakly on position, though the equivalent widths of all emission lines seem to decrease away from the Galactic center (green line, right panel).

sity and structure unresolved by IRAS introduces a bias, nearly independent of wavelength, which we estimate to be a factor of 2.1 ± 0.4 . We demonstrate both of these results below.

4.1. Measurement Errors

The errors in our correlation spectra are derived from fits to about 90,000 intensities over the full sky. The sky spectra are first interpolated onto a common wavelength array, introducing a correlation between the intensities at neighboring wavelengths. The difference between the intensities I_λ at wavelength elements i and $i + 1$ ranges from $\frac{1}{2}$ to 1 time(s) its value without interpolation (a factor of $\frac{1}{2}$ corresponds to the original and final arrays being offset by half a wavelength increment). Because the offsets between the original and resampled wavelengths are random, the denominator of this fraction is uniformly distributed between 1 and 2, which reduces the average difference $\langle I_{i+1} - I_i \rangle$ by a factor of

$$\left\langle \frac{(I_{i+1} - I_i)_{\text{orig}}}{(I_{i+1} - I_i)_{\text{interp}}} \right\rangle = \int_1^2 \frac{dt}{t} = \ln 2. \quad (10)$$

This correlation will propagate through the individual interpolated spectra to the α_λ . If the element-to-element variations in α_λ are dominated by Gaussian measurement errors, we expect the quantities

$$\Delta\alpha_i \equiv \frac{\alpha_{i+1} - \alpha_i}{(\ln 2) \sqrt{\sigma_{i+1}^2 + \sigma_i^2}} \quad (11)$$

to be normally distributed with unit variance; α_i and σ_i^2 are given by Equations (5) and (6), respectively.

Figure 7 shows that the distribution of $\Delta\alpha_i$ as defined by Equation (11) is exceedingly well-fit by a normal distribution with zero mean and unit variance. We have

masked 67 of the 3999 normalized intensity differences which lie near the emission lines $H\alpha$, $H\beta$, $[\text{N II}] \lambda 6550$, $[\text{N II}] \lambda 6585$, $[\text{S II}] \lambda 6718$, and $[\text{S II}] \lambda 6733$. At these wavelengths, real spectral features contribute to the element-to-element variation in α_λ . Even with 3932 values of $\Delta\alpha_i$ across the wavelength range, a Kolmogorov-Smirnov (K-S) test only detects deviation from Gaussianity with 83% confidence.

The exceptional agreement shown in Figure 7 gives us confidence that our measured errors are reliable and independent (up to a factor of $\ln 2$ from interpolating). We have verified that coadding neighboring wavelength elements, as we did to smooth the continuum in Section 3.1 and to compute the equivalent widths of lines in Section 3.2, does not affect the statistical properties of the errors.

4.2. Biases in the Correlation Spectra

Our model of the correlation between 100 μm intensity and optical intensity (Equation (2)) includes errors in the SDSS sky fiber residuals, but neglects (unknown) errors in the 100 μm intensity at each fiber's location and structure unresolved by IRAS. This introduces a wavelength-independent bias to our recovered spectra (Figures 3 - 6). We first derive an expression for the bias and then estimate its value.

The correlation spectrum α_λ (Equation (2)) is derived from a χ^2 minimization. The maximum likelihood values of the α_λ and their variances are given by Equations (5) and (6), with $x_{j,p}$ and $y_{\lambda,j,p}$ defined in Equations (3) and (4); $x_{j,p}$ is the excess 100 μm intensity in fiber j relative to the average on plate p , $y_{\lambda,j,p}$ is the residual sky fiber intensity λI_λ at wavelength λ in fiber j on plate p , and $\sigma_{\lambda,j,p}^2$ is its variance as estimated by the SDSS pipeline. We let $\xi_{j,p}$ denote the true excess 100 μm emission at sky fiber j on plate p , so that the measurement error,




FIG. 7.— Normalized difference between α_λ at successive wavelength elements (Equation (11)) in the full sky correlation spectrum, masking wavelengths near emission lines. The factor of $\ln 2$ arises from the fact that we interpolate all spectra onto a common wavelength range, which introduces a correlation between neighboring wavelength elements (see Section 4.1). There is no other scaling. The resulting distribution of 3932 points is very nearly Gaussian; a K-S test rejects the consistency of the two distributions with just 83% confidence.

including the effects of unresolved structure, is $\delta_{j,p} = x_{j,p} - \xi_{j,p}$.

Assuming the model in Equation (2) to be correct, we may write $y_{\lambda,j,p}$ as $\alpha_\lambda \xi_{j,p} + \varepsilon_{\lambda,j,p}$, with $\varepsilon_{\lambda,j,p}$ representing the measurement error in the sky fiber intensity. We further assume the error terms $\varepsilon_{\lambda,j,p}$ and $\delta_{j,p}$ to be uncorrelated with zero mean and invoke the large number of sky fibers (nearly 10^5) to neglect sums of δ and ε . The first factor in Equation (5) becomes

$$\sum_{j,p} \frac{(y_{\lambda,j,p})(x_{j,p})}{\sigma_{\lambda,y,j,p}^2} \approx \alpha_\lambda \sum_{j,p} \frac{\xi_{j,p}^2}{\sigma_{\lambda,y,j,p}^2}. \quad (12)$$

This is the same value we would measure with $\delta_{j,p} = 0$. Noting that the mean $100 \mu\text{m}$ excess, $\langle x_{j,p} \rangle$, is zero by construction, our estimate of α_λ is biased low by a factor

$$\frac{\alpha_\lambda(x, y)}{\alpha_\lambda(\xi, y)} \approx \left(\sum_{j,p} \frac{\xi_{j,p}^2}{\sigma_{\lambda,y,j,p}^2} \right) \left(\sum_{j,p} \frac{x_{j,p}^2}{\sigma_{\lambda,y,j,p}^2} \right)^{-1} \approx \frac{\sigma_\xi^2}{\sigma_x^2}, \quad (13)$$

where σ_x^2 is the observed variance in x and $\sigma_\xi^2 \approx \sigma_x^2 - \langle \delta^2 \rangle$ would be its value with no measurement errors. Thus,

$$\frac{\alpha_\lambda(x, y)}{\alpha_\lambda(\xi, y)} \approx 1 - \frac{\langle \delta^2 \rangle}{\sigma_x^2}. \quad (14)$$

Note that this bias is independent of wavelength and of the errors in the sky fiber intensities. We have empirically verified the latter by adding noise to the intensities; we recover the same correlation spectra to within the errors.

4.3. Calibrating the Correlation Spectra

An unbiased estimator for α_λ , for example a likelihood function of the form

$$\mathcal{L}(x_j, y_{\lambda,j} | \alpha_\lambda) = \int \mathcal{L}(x_j | x) \mathcal{L}(y_{\lambda,j} | x, \alpha_\lambda) \mathcal{L}(x) dx, \quad (15)$$

would remove all calibration issues. The SDSS pipeline provides an estimate of $\mathcal{L}(y_{\lambda,j} | x, \alpha_\lambda)$, the likelihood of measuring sky fiber residual y_λ in sky fiber j given a correlation spectrum α_λ and true $100 \mu\text{m}$ intensity x . Unfortunately, we have almost no information on the errors in the $100 \mu\text{m}$ intensity to estimate $\mathcal{L}(x_j | x)$, the likelihood of measuring excess $100 \mu\text{m}$ intensity x_j given a true value x , and can only guess at a prior, $\mathcal{L}(x)$, from the $100 \mu\text{m}$ map itself.

Because of these difficulties, we use two alternative and independent approaches. We first construct an estimator that we expect to be asymptotically unbiased, assuming the error in residual $100 \mu\text{m}$ intensity to depend weakly on the true residual value at a fiber position. This is a reasonable assumption, particularly because we are subtracting the mean $100 \mu\text{m}$ emission over a plate; an x -value of zero does *not* correspond to zero intensity. We then show the results of our radiative transfer calculations assuming ZDA04 dust and a plane-parallel galaxy.

Under the assumption of uniform errors in the residual $100 \mu\text{m}$ intensity, Equation (14) shows that restricting the sample to fibers with large $|x|$ (and therefore large σ_x^2) will give an asymptotically unbiased estimate. We therefore recalculate α_λ between 6600 and 6700 Å using only the fibers with $|x| > |x|_{\min}$. Figure 8 shows α_λ as a function of $|x|_{\min}$, calculated by varying $|x|_{\min}$ from 0 to 4 MJy sr⁻¹; as expected, α_λ increases with $|x|_{\min}$. Figure 8 suggests a bias factor of at least ~ 1.7 –2. This is also supported by our radiative transfer model, discussed in detail in Section 5.1, which suggests a bias of ≈ 1.7 –2.4. We conservatively adopt a bias factor of 2.1 ± 0.4 , indicated by the shaded region of Figure 8, to calibrate our correlation spectrum.

5. DISCUSSION

5.1. Scattered Light and Dust Models

Many features of the DGL – the 4000 Å break, broad Mg + Fe b absorption, and a much bluer continuum than that of stars with these spectral features – support the hypothesis that the DGL is dominated by scattered starlight. This is hardly surprising, as the spectrum was derived by correlating residual optical intensity with $100 \mu\text{m}$ intensity over small spatial scales. A simplified radiative transfer calculation confirms that our spectrum is consistent with dust scattering and allows us to use the DGL to discriminate between dust models.

Our model of the DGL uses an infinite plane-parallel galaxy with a Gaussian vertical distribution of dust,

$$\rho_{\text{dust}} \propto \exp \left[-\frac{z^2}{2\sigma^2} \right], \quad (16)$$

with $\sigma = 250$ pc (Malhotra 1995; Nakanishi & Sofue 2003). We approximate the stellar distribution as the sum of two exponential distributions with scale heights of 300 pc and 1350 pc (Binney & Merrifield 1998;




FIG. 8.— The correlation spectrum α_λ between 6600 and 6700 Å using only points with large excess 100 μm intensities. The red curve indicates the number of sky fibers satisfying each 100 μm cutoff. If the errors on residual 100 μm intensity depend weakly on fiber position, this estimator will be asymptotically unbiased. We overplot several single-scattering radiative transfer models, which we describe in detail in Section 5.1. All assume ZDA04 dust, but with different dust densities and different ISRFs. We conservatively adopt the bias factor of 2.1 ± 0.4 indicated by the orange shading to calibrate our correlation spectrum.

Gilmore & Reid 1983). The 300 pc component dominates the distribution with about 90% of the stars.

We use two estimates of the stellar emission spectrum:

1. A model that reproduces the local ISRF of Mathis et al. (1983), hereafter MMP83, in the mid-plane, and
2. A stellar population synthesis model from BC03, with solar metallicity and an exponential star formation history over 12 Gyr.

The MMP83 ISRF for $\lambda > 2460$ Å is approximated as a sum of three dilute blackbodies of $T = 3000$, 4000, and 7000 K, with dilution coefficients $W = 5 \times 10^{-13}$, 1.65×10^{-13} , and 10^{-14} , respectively. We compute the attenuation of the stellar emission using Equation (A3), setting the *attenuated* spectrum equal to the local ISRF. In this way, we “de-redden” the MMP83 ISRF to obtain the stellar source spectrum.

We show all of the ISRF spectra in Figure 9. The peak of the de-reddened MMP83 λF_λ is at about 8000 Å, corresponding to a temperature of ~ 4600 K. The spectrum is similar to a BC03 model with a 5 Gyr star formation timescale, though MMP83 has less UV emission. This is probably a result of the relative lack of early O stars in the Solar neighborhood and of our model’s neglect of extinction from dust in a young star’s birth cloud (Charlot & Fall 2000). The UV discrepancy becomes much more serious for BC03 models with more extended star formation histories. This UV excess significantly increases the dust heating and decreases the ratio of scattering in the optical to emission in the far-infrared. For a star formation timescale of 5 Gyr, this is a ~ 30 -40% effect relative to the de-reddened MMP83 model (see Figures 8 and 10).

Theoretical and observational estimates of the local star formation history favor roughly constant star formation rates over ~ 10 Gyr (Hernández et al. 2001;




FIG. 9.— Comparison of the MMP83 ISRF, de-reddened using our radiative transfer models, and BC03 models with solar metallicity and exponential star formation histories over 12 Gyr. The BC03 models have significant excesses in the UV that increase dust heating and reduce the ratio of the optical scattering to far-infrared emission.

Cignoni et al. 2006). Such models do not agree with our measured spectrum of the DGL and, using our radiative transfer model, would produce a very different ISRF from MMP83. This may indicate that a substantial fraction of the illuminating starlight originates relatively far from the Solar neighborhood, it may be a result of our simplified radiative transfer, or it may indicate that stars in the Solar neighborhood are generally older than is currently thought. The spatial variation of the DGL (Figure 4) shows that the geometry of the radiative transfer problem is likely to be important. A more detailed model of the Galaxy could better constrain the average stellar source spectrum.

Our radiative transfer calculations neglect multiple scatterings. This is a good approximation at high Galactic latitude where the optical depths are low. It becomes poor near the midplane, but SDSS has very little sky coverage near the Galactic plane (Figure 2). We assume all absorbed starlight to be reradiated isotropically in the infrared and use a Henyey-Greenstein phase function for scattering in the optical. We use the dust model of Draine & Li (2007) to convert total infrared power to IRAS 100 μm bandpass power, with

$$(\nu I_\nu)_{100 \mu\text{m}} = (0.52 \pm 0.05) I_{\text{TIR}}. \quad (17)$$

The central value corresponds to their model with an incident starlight intensity 80% of that in the Solar neighborhood, which Draine & Li (2007) found to provide the best fit to the average far-infrared spectrum measured by Finkbeiner et al. (1999). The confidence interval in Equation (17) includes models from 0.5 to 1.5 times the local starlight intensity. Once a dust model is specified, with wavelength-dependent cross-sections, albedos, and anisotropy parameters, our model for the scattered light




FIG. 10.— Single-scattering radiative transfer calculations assuming a plane-parallel galaxy. The results are much more sensitive to the assumed dust model than to the parameters of the galaxy model. The WD01 model produces more scattering at longer wavelengths than the ZDA04 model because of an excess of large dust grains.

spectrum has no free parameters. We derive the relevant equations in the Appendix and evaluate them numerically.

Figure 10 shows the results of our calculations to be remarkably insensitive to the details of the galaxy modeling (other than the assumed stellar source spectrum). Though not shown, models with a single exponential distribution of stars and with an exponential, rather than Gaussian, dust distribution, are nearly indistinguishable from the present models. For $\csc|b| > 1.4$ ($|b| < 45^\circ$) the H I has $N_{\text{H}} \approx 2.9 \times 10^{20} \csc|b| \text{ cm}^{-2}$ (Dickey et al. 1978) falling below this relation for $b > 45^\circ$. For $E(B - V)/N_{\text{H}} = 5.8 \times 10^{21} \text{ cm}^2 \text{ mag}^{-1}$ and $R_V = A_V/E(B - V) = 3.1$, we would then expect $\tau_V = 0.17 \csc|b|$ for $|b| < 45^\circ$, and lower values at $b > 45^\circ$. We use $\tau_V = 0.15 \csc|b|$ for our fiducial distribution.

Figure 11 shows that the optical depths at the SDSS sky fibers, computed using the SFD estimates of $E(B - V)$ and assuming $R_V = 3.1$ dust, are roughly bracketed by $\tau_V = 0.05 \csc|b|$ and $\tau_V = 0.15 \csc|b|$. This suggests that the sky fiber locations were slightly biased toward lower-than-average H I column densities.

While the details of the model galaxy have little effect on the predicted spectrum, the dust model matters a great deal. The WD01 model appears to have too many large grains, giving too much scattering at long wavelengths. The size distribution of ZDA04 brings our model into much better agreement with the data (see Figure 3). While our model predicts the scattering spectrum to be a very weak function of Galactic latitude, variations in the dust properties, geometry, or illuminating starlight could result in much larger differences. Possible spatial variations in $100 \mu\text{m}$ errors and small-scale structure (Section




FIG. 11.— Optical depths at each sky fiber computed using the SFD estimates of $E(B - V)$ and assuming $R_V = 3.1$ dust. The dust distribution is roughly bracketed by $\tau_V = 0.05 \csc|b|$ and $\tau_V = 0.15 \csc|b|$; these two models predict nearly the same spectrum of scattered light.

4.2) further complicate the interpretation of the spatial variations seen in Figure 4.

5.2. Line Emission

On the evidence presented above, we can be confident that the continuum of the correlation spectrum α_λ consists primarily of scattered starlight. It is more difficult to show that the line emission is scattered rather than from ionized gas physically associated with the dust. The strongest piece of evidence is that the equivalent width of H α in the correlation spectrum, $12.5 \pm 0.5 \text{ Å}$, matches its measured value of 11 Å in the local ISRF (Section 3.2). Because scattering will preserve the equivalent width of H α in the ISRF incident on the dust, a stronger H α line would have indicated an additional, nearly continuum-free component seen in direct emission and correlating with $100 \mu\text{m}$ intensity.

If the emission lines observed in Section 3.2 are observed mostly or entirely in reflection, they provide a probe of the average physics of the nearby ISM. As discussed in Section 3.2, the strength of the [N II] and [S II] lines, combined with the weakness of the [O III] lines, indicate relatively few photons with $h\nu > 24.6 \text{ eV}$. This is probably due to the lack of early-type O stars in the Solar neighborhood. The observed strengths of the collisionally-excited lines relative to recombination lines of similar wavelength are:

$$\frac{[\text{S II}](6718 + 6733)}{\text{H}\alpha 6565} = 0.80 \pm 0.05, \quad (18)$$

$$\frac{[\text{N II}](6550 + 6585)}{\text{H}\alpha 6565} = 0.72 \pm 0.06, \text{ and} \quad (19)$$

$$\frac{[\text{O III}](4960 + 5008)}{\text{H}\beta 4863} = 0.58 \pm 0.21. \quad (20)$$




FIG. 12.— Upper panel: $[N II](6550+6585)/H\alpha$ vs. T . Curves are predicted values for indicated values of $N II/H II$ relative to $(N/H)_{\odot}$. If $0.7 < (N II/H II)/(O/N)_{\odot} < 1$, then the observed line ratio 0.72 ± 0.06 requires $7180 < T < 8400 K$. Middle panel: $[S II](6718+6733)/H\alpha$ vs. electron temperature T . Curves are predicted values for indicated values of $S II/H II$ relative to $(S/H)_{\odot}$. The observed line ratio 0.80 ± 0.05 and $0.7 < (S II/H II)/(O/H)_{\odot} < 1$ limits the temperature to $7180 < T < 8330 K$. Lower panel: $[O III](4960+5008)/H\beta$ vs. T . Curves are predicted values for indicated values of $O III/H II$ relative to $(O/H)_{\odot}$. The observed line ratio 0.58 ± 0.21 and the allowed temperature range $7180 < T < 8330 K$ implies that $O III/H II$ is between $0.039 \times$ and $0.15 \times (O/H)_{\odot}$. The dot in each panel indicates our best estimate: $T = 7700 K$, $N II/H II = 0.85(N/H)_{\odot}$, $S II/H II = 0.83(S/H)_{\odot}$, and $O III/H II = 0.08(O/H)_{\odot}$.

These allow us to estimate the temperature of the ISM where the lines originate and the state of ionization of S, N, and O. Figure 12 shows predicted line ratios as functions of electron temperature T , calculated using $H\alpha$ and $H\beta$ emissivities from Draine (2011), collision strengths for N II from Hudson & Bell (2005), for S II from Tayal & Zatsarinny (2010), and for O III from Aggarwal & Keenan (1999). A density $n_e = 10^2 \text{ cm}^{-3}$ was assumed; the results are insensitive to n_e provided $n_e \lesssim 10^3 \text{ cm}^{-3}$.

Nitrogen is not depleted in the ISM, and the first and second ionization potentials (14.0 and 29.6 eV) lead us to expect $N II/H II$ to be close to the solar abundance $(N/H)_{\odot} = 7.4 \times 10^{-5}$ (Asplund et al. 2009). If we assume $0.7 < (N II/H II)/(N/H)_{\odot} < 1$ we see from Figure 12 that the observed $[N II](6550+6565)/H\alpha$ ratio allows only temperatures $7180 < T < 8400 K$.

Sulfur is not expected to be depleted in H II regions or the diffuse ISM. With an ionization potential of 23.38

eV for $S II \rightarrow S III$, S II will be the dominant ionization stage in H II regions where He is neutral. If we assume $S II/H II$ to be between 0.7 and $1.0 \times$ the solar abundance $(S/H)_{\odot} = 1.45 \times 10^{-5}$ (Asplund et al. 2009), then we see from Figure 12 that the observed $[S II(6718+6733)]/H\alpha$ requires $7180 < T < 8330 K$.

Oxygen is only slightly depleted in the diffuse ISM, with $\sim 20\%$ of the O resident in silicates. The second ionization potential of oxygen is 35.1 eV, and therefore O III will be present only when He is ionized. A star of spectral type O8 or earlier is required for the He ionization zone to account for more than 50% of the mass in the H II region (Draine 2011). As seen in Table 2, the nearest such stars are 15 Mon (O7V, $d = 309^{+60}_{-43}$ pc) and ζ Pup (O4I, $d = 335^{+12}_{-11}$ pc). The reflected light in the DGL is expected to originate mainly within a few hundred pc of the Sun, and therefore the contribution from H II regions should be dominated by H II regions where He is neutral and O is singly ionized. Figure 12 shows that the observed strength of $[O III](4960+5008)/H\beta$ is consistent with O III/H II between 0.039 and 0.15 of $(O/H)_{\odot} = 5.4 \times 10^{-4}$ (Asplund et al. 2009). The observed [O III] emission can be reproduced by emission from H II regions with $T \approx 7700 K$ and $O III/H II \approx 0.08 \times (O/H)_{\odot}$.

He I 5877 is not detected, with a $3\text{-}\sigma$ upper limit $He I 5877/H\alpha < 0.24$; for $T \approx 7700 K$, this corresponds to $He II/H II < 0.48$, which is not a useful constraint.

The emission lines measured in the correlation spectrum α_{λ} are therefore consistent with the line ratios expected for H II regions within ~ 400 pc of the Sun, for electron temperature $T \approx 7700 \pm 250 K$. These temperatures and ionization states are also consistent with the WIM (Madsen et al. 2006), which we expect to contribute a significant fraction of the line emission, particularly at high latitudes.

5.3. The Fraction of Scattered $H\alpha$ at High Latitude

The previous section argued that the $H\alpha$ in our correlation spectrum is scattered. We can then use the calibrated correlation spectra to estimate the fraction of the observed $H\alpha$ at high Galactic latitude that is scattered light. We simply integrate the correlation spectrum over the $H\alpha$ line and multiply by the ratio of the total $100 \mu m$ to $H\alpha$ emission over the high latitude sky fibers.

From Figures 8 and 10, we take the ratio $\lambda I_{\lambda}(6600 \text{ \AA})/\nu I_{\nu}(100 \mu m)$ to be 0.38 ± 0.07 . This continuum value allows us to convert an $H\alpha$ equivalent width in our correlation spectrum into an intensity relative to the $100 \mu m$ intensity. We then have

$$\frac{I(H\alpha)}{R} \approx (0.090 \pm 0.017) \times \left(\frac{W_{\lambda}(H\alpha)}{12.5 \text{ \AA}} \right) \times \left(\frac{I_{\nu}(100 \mu m)}{\text{MJy sr}^{-1}} \right), \quad (21)$$

where the $H\alpha$ intensity $I(H\alpha)$ is in Rayleighs² and we recall (Table 1) that $W_{\lambda}(H\alpha) \approx 12.5 \text{ \AA}$ in our correlation spectrum. Our result (21) is close to the value $I(H\alpha)/R = (0.129 \pm 0.015)(I_{\nu}(100 \mu m)/\text{MJy sr}^{-1})$ found recently by Witt et al. (2010).

Averaging the $100 \mu m$ intensity from SFD and the $H\alpha$ intensity from Finkbeiner (2003) over all regions with

² $1 R = \frac{10^6}{4\pi} \text{ photons cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$

$b > 60^\circ$ gives a ratio

$$\left\langle \frac{I(\text{H}\alpha)}{R} \right\rangle = 0.47 \left\langle \frac{I_\nu(100 \mu\text{m})}{\text{MJy sr}^{-1}} \right\rangle. \quad (22)$$

The ratio of the coefficients in (21,22) gives a scattered fraction of H α at high latitudes of $0.090/0.47 = 0.19$. We add the statistical error of 5% in the H α equivalent width and our conservative estimate of the uncertainty in the calibration shown in Figure 8 in quadrature, giving a scattered H α fraction at high latitude of

$$\frac{\text{scattered H}\alpha}{\text{total H}\alpha} = 0.19 \pm 0.04. \quad (23)$$

Witt et al. (2010) produced a scatter diagram of the scattered H α fraction (their Fig. 6). The centroid of their distribution appears to be close to our measured value 0.19 ± 0.04 .

Wood & Reynolds (1999) estimated that 5–20% of the H α at high latitudes would be scattered light from H II regions. In a theoretical study of the emission spectrum of the diffuse H α , Dong & Draine (2011) concluded that the observed line ratios, including the low ratio of radio free-free to H α , could be understood if $\sim 20\%$ of the diffuse H α is actually reflected from dust, rather than emission from recombining gas in that direction. All of these results are consistent with our value 0.19 ± 0.04 .

We expect the scattered H α to originate both from H II regions around young stars and from the WIM. Our line ratios seem to be more consistent with those of the WIM than with classical H II regions (Madsen et al. 2006), which may indicate that the WIM dominates the line emission in our correlation spectra. However, this could also be due to the lack of nearby H II regions illuminated by early O stars. Our line ratios do seem to be compatible with those in the ϕ Per H II region, which is illuminated by a B0.5+sdO system (Fig. 12 of Madsen et al. 2006).

5.4. Dust Luminescence

We find no evidence of the Extended Red Emission (ERE) observed in some reflection nebulae (e.g., NGC 7023) as a broad emission excess peaking near 7000 Å, with a FWHM ~ 1500 Å (Witt & Vijn 2004, and references therein); a similar broad excess is reported to be present in the diffuse ISM (Gordon et al. 1998; Szomoru & Guhathakurta 1998; Witt et al. 2008). The spectrum of light reflected by cirrus clouds was measured by Szomoru & Guhathakurta (1998), who reported a broad peak in λI_λ near 6500 Å that was attributed to luminescence, on the grounds that scattering is insufficient. Our study also finds λI_λ peaking near 6500 Å (see Figure 3) but the observed spectrum appears to be consistent with what is expected for scattering. Indeed, our scattering models tend to *over*-predict the DGL around 7000–9000 Å (Figure 3), depending on the adopted dust size distribution.

If we posit that the slight excess emission between 5500 Å and 7500 Å over our prediction with the ZDA04 dust model (see Fig. 3) is an upper limit on the ERE, we infer that dust luminescence accounts for no more than 10% of the DGL in this wavelength range and less elsewhere. After calibrating the spectra and relating $I_\nu(100 \mu\text{m})$ to

the total IR power using Eq. (17), this implies a ratio of ERE to infrared power of $\lesssim 0.4\%$, or a ratio of ERE to scattered power of $\lesssim 1\%$.

The ERE will be a function of the intensity of illuminating UV light and the dust column density. It is possible that our technique, which requires spatial variations of the ERE correlated with 100 μm intensity over a $\sim 1^\circ$ scale, is ill-suited to its detection. If the UV ISRF is particularly weak or spatially variable on small scales, it could at least partially explain the lack of ERE in our correlation spectrum. We have therefore performed two tests.

First, we made our optical depth cutoff $A_V \lesssim 0.5$ more stringent to restrict the calculation to regions optically thin to UV photons. Reducing this cutoff by a factor of 2 had a negligible effect on the correlation spectrum. Unfortunately, when we reduced it further, the main stellar spectral features disappeared and new features similar to those in the high latitude spectrum of Figure 4 appeared. At such low levels of 100 μm emission, we expect extragalactic contamination to be significant.

We have also tried to estimate empirically how much of the small-scale variation (over $\sim 1^\circ$) in 100 μm emission is due to variations in the illuminating starlight and how much is due to variations in the dust column density. We recalculated the correlation spectrum, correlating the SDSS intensities against dust optical depth rather than 100 μm intensity. Our recovered correlation spectrum was nearly identical. Unfortunately, the resolution of the SFD temperature map used to convert from 100 μm intensity to optical depth, 0.7° , is too coarse to make this test conclusive.

While we find no evidence of hidden ERE in our correlation spectrum, a more thorough investigation may require instruments that can more directly probe the dust-scattered optical spectrum.

6. SUMMARY AND CONCLUSIONS

In this paper, we have measured the spectrum of the scattered component of the DGL using 90,000 calibration spectra from SDSS by correlating against 100 μm intensity measured by COBE and IRAS and reduced by SFD. The correlation spectrum of the DGL is consistent with scattering of photons emitted by stars and by ionized gas. Its continuum and lines show interesting variations across the sky, which could be due to differences in the illuminating starlight, in the dust properties, or structure in the Galaxy. Our spectrum is not calibrated because the correlation neglects (unknown) errors and small-scale structure in the 100 μm intensity. However, an asymptotically unbiased estimator agrees well with simplified radiative transfer calculations and allows us to calibrate the average spectrum of the DGL. In calibrating the data, we also provide an estimate of the errors and small-scale structure in the SFD 100 μm map.

With a simplified radiative transfer calculation assuming a plane-parallel galaxy, we show that the spectrum of the DGL can discriminate between dust models. The ZDA04 dust model is preferred to the WD01 model, which has more large grains and scatters too much red light. The ZDA04 model fits the data very well without requiring dust luminescence as a source of ERE. Indeed, our radiative transfer calculations tend to over-predict scattering redward of 7000 Å.

Our measurement of $H\alpha$ in the DGL allows us to indirectly measure the fraction of scattered $H\alpha$ at high Galactic latitude. We find an average value of 0.19 ± 0.04 , consistent with the results of Wood & Reynolds (1999), Witt et al. (2010) and the value inferred by Dong & Draine (2011). The line emission also allows us to constrain the properties of local H II regions. The lack of emission from species with high ionization energies is consistent with the lack of nearby early O stars, and the average scattered spectrum is consistent with gas at ~ 7500 K.

Our results are the product of the extraordinary SDSS, the only dataset of its kind. It would be possible, though difficult, to incorporate the new Baryon Oscillations Spectroscopic Survey spectra (BOSS, Eisenstein et al. 2011) into the analysis. The new fibers are smaller ($2''$ instead of $3''$), losing far more light and making spectrophotometric calibration much more difficult. The DGL is a sufficiently weak signal that it requires thousands of blank sky spectra to extract useful results. These spectra also need to be in regions with significant dust columns, regions that tend to be avoided by large extragalactic surveys. The SDSS spectra will likely remain the best tool for analyzing the DGL for some time.

While the optical data are unlikely to improve in the near future, WISE and AKARI may allow significant improvements to the infrared map. Our results indicate

that the measurement errors and small-scale structure in the SFD $100 \mu\text{m}$ map are nearly as large as the $\sim 1^\circ$ variations to which our correlations are sensitive. By combining high resolution infrared data with the SFD maps, we could improve the quality of our DGL spectra and resulting constraints on Galactic stars and dust.

We wish to thank Michael Strauss, Dan Maoz, Yasushi Suto, and Steve Bickerton for many helpful conversations, and Michael Strauss for numerous suggestions that greatly improved the manuscript. We also wish to thank Jim Gunn and Robert Lupton for their help understanding the intricacies of SDSS data. This material is based upon work supported by the National Science Foundation Graduate Research Fellowship under Grant No. DGE-0646086. This research was supported in part by NSF grant AST-1008570.

Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS Web Site is <http://www.sdss.org/>.

APPENDIX

SCATTERING AND ABSORPTION IN A PLANE-PARALLEL EXPONENTIAL GALAXY

We test dust models by calculating scattered spectra in an infinite plane-parallel galaxy, using vertical distributions of stars and dust appropriate to the Solar neighborhood. The Sun is assumed to lie in the midplane, $z = 0$. We calculate the absorption (and hence, infrared emission) and scattering along each line-of-sight, neglecting multiple scatterings.

We denote the (wavelength-dependent) optical depth of dust above height z along a vertical line-of-sight as $\tau_\lambda(z)$,

$$\tau_\lambda(z) \equiv \int_z^\infty \sigma_{\text{ext}}(\lambda) \rho(z') dz' , \quad (\text{A1})$$

where σ_{ext} is the extinction cross-section and ρ is the dust density. Consider a sheet of stars with uniform surface power density Σ_λ at height z_s . The optical depth from an annulus of stars at a distance R from the grain's projected position in the stellar sheet to a grain at height z is

$$A_\lambda(z, z_s, R) \equiv |\tau_\lambda(z) - \tau_\lambda(z_s)| \frac{\sqrt{(z - z_s)^2 + R^2}}{|z - z_s|} . \quad (\text{A2})$$

The total flux density (neglecting multiply scattered photons) incident on a grain at height z is then

$$\begin{aligned} F_\lambda(z, z_s) &= \int_0^\infty 2\pi R dR \frac{\Sigma_\lambda \exp[-A_\lambda(z, z_s, R)]}{4\pi [(z - z_s)^2 + R^2]} \\ &= \left(\frac{\Sigma_\lambda}{2} \right) E_1(|\tau_\lambda(z) - \tau_\lambda(z_s)|) , \end{aligned} \quad (\text{A3})$$

where E_1 is the exponential integral and τ_λ is related to z by Equation (A1). The total reradiated intensity is an integral of the absorbed fraction of Equation (A3) over all wavelengths and stellar sheets. Defining ω_λ as the albedo, the total infrared intensity I_{TIR} from a sightline at Galactic latitude b is

$$I_{\text{TIR}}(b) = \frac{\csc|b|}{8\pi} \int_0^\infty (1 - \omega_\lambda) d\lambda \int_{-\infty}^\infty \frac{d\Sigma_\lambda}{dz_s} dz_s \int_0^{\tau_\lambda(0)} d\tau' E_1(|\tau' - \tau_\lambda(z_s)|) . \quad (\text{A4})$$

A dust model, like the Draine & Li (2007) model used in Equation (17), is needed to convert I_{TIR} into a specific intensity I_ν at infrared frequency ν .

Anisotropic scattering makes it more difficult to calculate the optical intensity. Consider an annulus of the stellar sheet centered directly below a dust grain, and let θ sweep out the annulus, with $\theta = 0$ pointed away from us. If the

dust grain is along a sightline at Galactic latitude b , the law of cosines gives the required scattering angle ξ as

$$\cos \xi = \frac{(z - z_s)^2 - Rz \cos \theta \cot b}{\sqrt{(z^2 \cot^2 b + (z - z_s)^2)(R^2 + (z - z_s)^2)}}. \quad (\text{A5})$$

The intensity of light scattered in our direction from a stellar sheet at height z_s is then given by

$$I_{\lambda, \text{sca}}(z, z_s, b) = \int_0^\infty R dR \int_0^{2\pi} d\theta \phi_\lambda(\cos \xi) \frac{\Sigma_\lambda \exp[-A_\lambda(z, z_s, R)]}{4\pi((z - z_s)^2 + R^2)}, \quad (\text{A6})$$

with ϕ_λ being the normalized phase function and A_λ defined by Equation (A2). We use a Henyey-Greenstein phase function for simplicity. If ϕ is constant, the θ integral is trivial and we recover Equation (A3) modulo a factor of 4π . Scattered light is further attenuated by dust along the line-of-sight. Neglecting multiple scatterings, we finally have

$$I_{\lambda, \text{sca}}(z_s, b) = \omega_\lambda \csc |b| \int_0^{\tau_\lambda(0)} \exp[-\csc |b|(\tau_\lambda(0) - \tau')] d\tau' \int_0^\infty R dR \int_0^{2\pi} d\theta \phi_\lambda(\cos \xi) \frac{\Sigma_\lambda \exp[-A_\lambda(z, z_s, R)]}{4\pi((z - z_s)^2 + R^2)}, \quad (\text{A7})$$

where z and τ_λ are related by Equation (A1) and A_λ is defined by Equation (A2). As for Equation (A4), we integrate Equation (A7) over z_s . We do not integrate over wavelength; the spectrum of $I_{\lambda, \text{sca}}$ is shown in Figure 3 normalized by the infrared intensity νI_ν at $100 \mu\text{m}$, which is related to I_{TIR} by Equation (17). The ratio $I_{\lambda, \text{sca}}/I_{\text{TIR}}$ depends very weakly on Galactic latitude b , the normalization of the optical depth, and the details of the stellar and dust distributions. Once an ISRF and dust model are supplied, there are no other free parameters.

REFERENCES

- Abazajian, K. N., Adelman-McCarthy, J. K., Agüeros, M. A., et al. 2009, *ApJS*, 182, 543
- Adelman-McCarthy, J. K., Agüeros, M. A., Allam, S. S., et al. 2008, *ApJS*, 175, 297
- Adelman-McCarthy, J. K., Agüeros, M. A., Allam, S. S., et al. 2006, *ApJS*, 162, 38
- Aggarwal, K. M., & Keenan, F. P. 1999, *ApJS*, 123, 311
- Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, *ARA&A*, 47, 481
- Binney, J., & Merrifield, M. 1998, *Galactic Astronomy* (Princeton, NJ: Princeton University Press)
- Boggess, N. W., Mather, J. C., Weiss, R., et al. 1992, *ApJ*, 397, 420
- Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000
- Charlot, S., & Fall, S. M. 2000, *ApJ*, 539, 718
- Cignoni, M., Degl'Innocenti, S., Prada Moroni, P. G., & Shore, S. N. 2006, *A&A*, 459, 783
- Dennison, B., Simonetti, J. H., & Topasna, G. A. 1998, *PASA*, 15, 147
- Dickey, J. M., Terzian, Y., & Salpeter, E. E. 1978, *ApJS*, 36, 77
- Dong, R., & Draine, B. T. 2011, *ApJ*, 727, 35
- Draine, B. T. 2011, *Physics of the Interstellar and Intergalactic Medium* (Princeton, NJ: Princeton University Press)
- Draine, B. T., & Li, A. 2007, *ApJ*, 657, 810
- Eisenstein, D. J., Weinberg, D. H., Agol, E., et al. 2011, *AJ*, 142, 72
- Elsässer, H., & Haug, U. 1960, *ZAp*, 50, 121
- Elvey, C. T., & Roach, F. E. 1937, *ApJ*, 85, 213
- Finkbeiner, D. P. 2003, *ApJS*, 146, 407
- Finkbeiner, D. P., Davis, M., & Schlegel, D. J. 1999, *ApJ*, 524, 867
- Frieman, J. A., Bassett, B., Becker, A., et al. 2008, *AJ*, 135, 338
- Gaustad, J. E., McCullough, P. R., Rosing, W., & Van Buren, D. 2001, *PASP*, 113, 1326
- Gilmore, G., & Reid, N. 1983, *MNRAS*, 202, 1025
- Gordon, K. D., Witt, A. N., & Friedmann, B. C. 1998, *ApJ*, 498, 522
- Henry, R. C. 1981, *ApJ*, 244, L69
- Hernández, X., Avila-Reese, V., & Firmani, C. 2001, *MNRAS*, 327, 329
- Hudson, C. E., & Bell, K. L. 2005, *A&A*, 430, 725
- Hurwitz, M., Bowyer, S., & Martin, C. 1991, *ApJ*, 372, 167
- Lillie, C. F., & Witt, A. N. 1976, *ApJ*, 208, 64
- Madsen, G. J., Reynolds, R. J., & Haffner, L. M. 2006, *ApJ*, 652, 401
- Maíz Apellániz, J., Alfaro, E. J., & Sota, A. 2008, *ArXiv e-prints*, 0804.2553
- Maíz-Apellániz, J., Walborn, N. R., Galué, H. Á., & Wei, L. H. 2004, *ApJS*, 151, 103
- Malhotra, S. 1995, *ApJ*, 448, 138
- Martin, C., Hurwitz, M., & Bowyer, S. 1990, *ApJ*, 354, 220
- Mathis, J. S., Mezger, P. G., & Panagia, N. 1983, *A&A*, 128, 212
- Morgan, D. H., Nandy, K., & Thompson, G. L. 1978, *MNRAS*, 185, 371
- Murthy, J., Henry, R. C., Feldman, P. D., & Tennyson, P. D. 1990, *A&A*, 231, 187
- Murthy, J., Henry, R. C., & Holberg, J. B. 1991, *ApJ*, 383, 198
- Nakanishi, H., & Sofue, Y. 2003, *PASJ*, 55, 191
- Neugebauer, G., Habing, H. J., van Duinen, R., et al. 1984, *ApJ*, 278, L1
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2011, *ArXiv e-prints*, 1101.2029
- Reynolds, R. J., Haffner, L. M., & Madsen, G. J. 2002, in *ASPC*, Vol. 282, *Galaxies: the Third Dimension*, ed. M. Rosada, L. Binette, & L. Arias, 31
- Sasseen, T. P., & Deharveng, J.-M. 1996, *ApJ*, 469, 691
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Seon, K.-I., Edelstein, J., Korpela, E., et al. 2011, *ApJS*, 196, 15
- Stoughton, C., Lupton, R. H., Bernardi, M., et al. 2002, *AJ*, 123, 485
- Szomoru, A., & Guhathakurta, P. 1998, *ApJ*, 494, L93
- Tayal, S. S., & Zatsarinny, O. 2010, *ApJS*, 188, 32
- Weingartner, J. C., & Draine, B. T. 2001, *ApJ*, 548, 296
- Witt, A. N., Gold, B., Barnes, F. S., III, et al. 2010, *ApJ*, 724, 1551
- Witt, A. N., Mandel, S., Sell, P. H., Dixon, T., & Vijh, U. P. 2008, *ApJ*, 679, 497
- Witt, A. N., & Vijh, U. P. 2004, in *ASPC 309, Astrophysics of Dust*, ed. A. N. Witt, G. C. Clayton, & B. T. Draine (San Francisco, CA: ASP), 115
- Wolstencroft, R. D., & Rose, L. J. 1966, *Nature*, 209, 388
- Wood, K., & Reynolds, R. J. 1999, *ApJ*, 525, 799
- Yahata, K., Yonehara, A., Suto, Y., et al. 2007, *PASJ*, 59, 205
- York, D. G., Adelman, J., Anderson, J. E., Jr., et al. 2000, *AJ*, 120, 1579
- Zubko, V., Dwek, E., & Arendt, R. G. 2004, *ApJS*, 152, 211
- Zvereva, A. M., Severnyi, A. B., Granitskii, L. V., et al. 1982, *A&A*, 116, 312